

BRIGHAM HEALTH  
BRIGHAM AND  
WOMEN'S HOSPITAL

## Can't-Miss Radiology Diagnoses

Jennifer W. Uyeda, MD  
Assistant Professor of Radiology  
Brigham and Women's Hospital  
Harvard Medical School

HARVARD MEDICAL SCHOOL  
TEACHING HOSPITAL

A KENNEDY MEDICAL GROUP PARTNERS  
HOSPITAL

BRIGHAM HEALTH  
BRIGHAM AND  
WOMEN'S HOSPITAL

HARVARD MEDICAL SCHOOL  
TEACHING HOSPITAL

## Financial Disclosures


- None

## Lecture Outline

- Over the next 50 minutes, the audience will be able to:
  - Interpret basic chest radiographs
  - Recognize and locate various tubes and lines
  - Identify “can’t miss radiology diagnoses” on plain x-ray and CT
  - Discuss various imaging protocols and considerations
  - Recognize the pros and cons of IV and PO contrast use in CT
  - Optimally triage renal failure and contrast allergy patients

## Part I: Interpreting chest radiographs





**BRIGHAM HEALTH**  
BRIGHAM AND  
WOMEN'S HOSPITAL

## Conventional Radiographs


- Four basic tissue densities
  1. Air
  2. Fat
  3. Tissue (water)
  4. Metal (bone)

#1

## Normal Chest Radiograph

Right paratracheal  
stripe: SVC

Right heart border:  
Right atrium



Trachea


Aortic knob and  
descending  
thoracic aorta

Left heart border:  
left ventricle



AP (L)

Opacity with air  
bronchograms



(L)





Left Upper Lobe Pneumonia: air bronchograms













## Atelectasis


Imaging findings:

Whiteout with volume loss-  
elevation of hemidiaphragm  
and ipsilateral tracheal &  
mediastinal shift



## Large Pleural Effusion

Imaging findings:

Whiteout with mass effect-  
contralateral tracheal &  
mediastinal shift




1200 mL serosanguineous fluid





“Opacities could represent atelectasis, pleural effusion, or pulmonary edema, however, airspace disease/pneumonia cannot be excluded.”





#2

CT has better  
contrast discrimination





Tree in bud



## "Tree-in-bud" opacities

- Irregular, nodular branching pattern
- Represent dilated and impacted centrilobular bronchioles
- Infection (endobronchial spread)
  - bacterial, fungal, MAC
- Airway disease
  - CF, bronchiectasis
- Allergic bronchopulmonary aspergillosis
- Aspiration






## “Ground Glass” Opacities

- Increased lung attenuation without obscuring underlying vessels
- Replacement of air in alveoli by
  - Fluid (pus, edema, hemorrhage),
  - Cells (tumor),
  - Fibrosis






#3

One View  
is  
No View




## Part II: Tubes and Lines





Q1. What is the most concerning finding on this CXR?


- A. Location of feeding tube
- B. Pulmonary edema
- C. Left pneumothorax
- D. Right lung pneumonia



Q1. What is the most concerning finding on this CXR?

- A. Location of feeding tube
- B. Pulmonary edema
- C. Left pneumothorax
- D. Right lung pneumonia






### A. Location of feeding tube

Dobhoff tube is incorrectly located in the right mainstem bronchus.


Correct position: esophageal with tip below the diaphragm.

Do NOT feed the patient!  
Reposition Dobhoff.




Q2. What is the next best step for this patient?

- A. SBO oral contrast pathway
- B. Exploratory laparotomy
- C. Barium enema
- D. Thoracic surgery consultation
- E. Urology consultation



Q2. What is the next best step for this patient?

- A. SBO oral contrast pathway
- B. Exploratory laparotomy
- C. Barium enema
- D. Thoracic surgery consultation**
- E. Urology consultation



**D. Thoracic surgery consultation**

Dobhoff tube is incorrectly located in the distal right lower lobe bronchus.

Correct position: esophageal with tip below the diaphragm.


Do NOT feed the patient!

Distal position of the Dobhoff may have perforated small bronchiole and could cause a tension pneumothorax upon removal


R IJ Swan Ganz catheter is  
incorrectly located in the  
distal right pulmonary artery.

Correct position: Proximal  
right pulmonary artery.

Retract catheter.




## Part III: Can't Miss Radiology Diagnoses



Q3. What is the most concerning finding on this CXR?


- A. Location of Port catheter
- B. Left pleural effusion
- C. Left pneumothorax
- D. Tension pneumothorax



Q3. What is the most concerning finding on this CXR?

- A. Location of Port catheter
- B. Left pleural effusion
- C. Left pneumothorax
- D. Tension pneumothorax





## D. Tension Pneumothorax


Imaging findings:

Pneumothorax with

Ipsilateral increased intercostal spaces

Contralateral mediastinal shift

Depression of diaphragm



## Left pneumothorax



Imaging findings:



Absent lung markings

Visceral pleural line

Peripheral space is lucent

Lung may be collapsed






## Subtle free subdiaphragmatic air


Identify liver & diaphragm  
Lucency under hemidiaphragm

UPRIGHT radiograph is helpful  
to identify antidependent air





- Upright CXR can depict intraperitoneal gas as a lucency under the diaphragm
- CT is more sensitive for the detection of pneumoperitoneum and CT also shows the site of perforation in up to 85% of actual perforation sites.




## Perforated Bowel: Peptic Ulcer


One of the most common causes of perforated bowel



## Free Intraperitoneal Air

- Abdominal emergency requiring surgical or percutaneous intervention
- Contained perforations can be managed conservatively
- Most common sources:
  1. Perforated diverticulitis
  2. Perforated peptic ulcers
  3. Others: perforated carcinoma, bowel ischemia
- Iatrogenic:
  - Post endoscopy, colonoscopy
  - Postoperative






## Rigler Sign

Imaging findings:

Air is seen on both sides of the bowel wall (intraluminal AND peritoneal side)





## Small Bowel Obstruction

Imaging findings:


Dilated small bowel loops  $>3\text{cm}$   
Air-fluid levels



## Small Bowel Obstruction

Imaging findings:

Dilated small bowel loops  $>3\text{cm}$   
Air-fluid levels




## BWH Adhesive SBO pathway




- Trial of conservative management of SBO
- Evidence
  - 80% SBO will resolve without surgery
  - High grade SBO can be safely managed non-operatively
  - Iodinated water soluble PO contrast that reaches colon within 24 hours indicates SBO will resolve without surgery
    - 97% sensitivity, 96% specificity
  - Current guidelines recommend that patients without resolution undergo surgery by 3-5 days
- Exclusion criteria:
  - Cancer, pregnancy, intra-abdominal infection

### Adhesive Small Bowel Obstruction Pathway









## Saddle Pulmonary Embolism

Imaging findings:

Filling defects in the pulmonary arteries





## Right Heart Strain in Pulmonary Emboli



Bowing of  
interventricular  
septum

RV: LV > 1:1





## Aortic Aneurysm Rupture


Break in wall of abdominal  
aortic aneurysm

I+: AAA with active  
extravasation


I-: focal discontinuity of  
calcification  
High attenuation hematoma



Aortic Aneurysm Rupture



Previous CTA



Subarachnoid  
hemorrhage



Subdural  
hemorrhage



Midline shift,  
herniation

## Part IV: Imaging Protocols and Considerations



## Imaging Tool Box

- Radiography/fluoroscopy
- Ultrasound (US)
- Computed tomography (CT)
- Magnetic resonance imaging (MR)
- Nuclear medicine (PET/CT, GI bleeding scan, HIDA, bone scan, etc)


BRIGHAM HEALTH  
BRIGHAM AND  
WOMEN'S HOSPITAL

CT ABDOMEN/PELVIS


Abd\_ABD -\*\*\* SCAN ABDOMEN ONLY NO PELVIS\*\* {74160}  
Abd\_Abd-\*\*\*SCAN PELVIS ONLY NO ABDOMEN\*\*\* {72193}  
Abd\_Abd, GU, Kidney, Renal Mass {74170}  
Abd\_Abd-Adrenal, Adrenal Mass {74150, 74160, 74170}  
Abd\_Abd-GU, CT Cystogram {72193}  
Abd\_Abd-Liver, Liver Mass {74170}  
Abd\_Abd-Pel, Abdomen/Pelvis {74176, 74177, 74178}  
Abd\_Abd-Pel, GI, Colonography, Diagnostic {74262}  
Abd\_Abd-Pel, GI, Colonography, Screening {74263}  
Abd\_Abd-Pel, GI, Enterography Dual Phase {74177}  
Abd\_Abd-Pel, GI, Enterography Single Phase {74177}  
Abd\_Abd-Pel, GU, Stone Protocol (L-) {74176}  
Abd\_Abd-Pel, GU, Urogram {74178;76377}  
Abd\_Abd-Pel, GU, Urogram, Under 35 {74178;76377}  
Abd\_Abd-Pel, Hernia {74178};{74176}  
Abd\_Abd-Pel, Hypervascular Tumor Staging {74177}  
Abd\_Abd-Pel, Low Dose Follow Up Known Ureteral Stone {74176}  
Abd\_Abd-Pel, Panc Dual Phase {74177}  
Abd\_Abd-Pel, Panc Three Phase {74177;76377}  
Abd\_Ch-Abd, Chest/Abdomen ONLY \*\*\*NO PELVIS\*\*\* {71250, 71260, 71270, 74150, 74160, 74170}  
Abd\_Ch-Abd-Pel, Chest/Abdomen/Pelvis {71250, 71260, 71270, 74176, 74177, 74178}  
Abd\_Ch-Abd-Pel, Hypervascular Tumor Staging-Chest/Abdomen/Pelvis {71260, 74177}  
Abd\_Nk\_Ch-Ab-Pel, Neck/Chest/Abdomen/Pelvis {70490, 70491, 70492, 71250, 71260, 71270, 74176, 74177, 74178}  
CH\_Research Only {71250}  
ER\_Mesenteric and GI Bleeding CTA {74174}  
ER\_Trauma (Panscan Plus): Head, Face, CTLS-spine, C/A/P {70450;70486;72125;72129;72132;71260;74177}  
ER\_Trauma (Panscan): Head, CTLS-spine, C/A/P {70450;72125;72129;72132;71260;74177}  
ER\_Trauma, A/P, LS-spine {74177;72132}  
ER\_Trauma, C/A/P, TLS-spine {71260;74177;72129;72132}  
ER\_Trauma, Head, C-spine, A/P, LS-spine {70460;72125;74177;72131}

MEDICAL SCHOOL  
HOSPITAL

## Pancreatic Neuroendocrine Tumor





Arterial Phase CT AP



Portal Venous Phase CT AP

## Bolus Tracking Technique





87 yo woman with HTN, CAD, diabetes, renal failure, dementia presents with severe and acute abdominal pain. Abdominal pain is maybe ?out of proportion to physical examination.

Now what?  
Do I order an imaging study?  
What study do I order?

US

MRE

IR/Angio

CT


CT I-


CTA

## ACR Appropriateness Criteria®

The ACR Appropriateness Criteria® (AC) are **evidence-based guidelines** to assist referring physicians and other providers in making the **most appropriate imaging or treatment decision for a specific clinical condition**. Employing these guidelines helps providers enhance quality of care and contribute to the most efficacious use of radiology.

<https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>





[All](#)
[Images](#)

About 258,000


ACR Appr  
<https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>  
The ACR Appr  
and other provi


Myelop  
Cardiac, T  
Rating Tab

Overvie  
ACR Appr  
Overview.  
More resul

**ACR Appropriateness Criteria**

The ACR Appropriateness Criteria® (AC) are evidence-based guidelines to assist referring physicians and other providers in making the most appropriate imaging or treatment decision for a specific clinical condition. Employing these guidelines helps providers enhance quality of care and contribute to the most efficacious use of radiology. [Learn more »](#)



 See the complete list of ACR AC topics and ratings tables »

[Browse Topics ↗](#)

<https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>



Gastrointestinal				
Topic Name	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Acute Nonlocalized Abdominal Pain	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Acute Pancreatitis	Narrative & Rating Table	Evidence Table		Appendix
Blunt Abdominal Trauma	Narrative & Rating Table	Evidence Table		Appendix
Chronic Liver Disease	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Colorectal Cancer Screening	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Crohn Disease	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Dysphagia	Narrative & Rating Table	Evidence Table		Appendix
Imaging of Mesenteric Ischemia	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Jaundice	Narrative & Rating Table	Evidence Table		Appendix
Left Lower Quadrant Pain — Suspected Diverticulitis	Narrative & Rating Table	Evidence Table		Appendix
Liver Lesion — Initial Characterization	Narrative & Rating Table	Evidence Table		Appendix
Nonvariceal Upper Gastrointestinal Bleeding	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Palpable Abdominal Mass	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Pretreatment Staging of Colorectal Cancer	Narrative & Rating Table	Evidence Table	Lit Search	Appendix
Right Lower Quadrant Pain — Suspected Appendicitis	Narrative & Rating Table	Evidence Table		Appendix

<https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>

American College of Radiology ACR Appropriateness Criteria® Imaging of Mesenteric Ischemia		
<b>Variant 1:</b> Suspected acute mesenteric ischemia. Initial imaging.		
Procedure	Appropriateness Category	Relative Radiation Level
CTA abdomen and pelvis with IV contrast	Usually Appropriate	☆☆☆
CT abdomen and pelvis with IV contrast	May Be Appropriate	☆☆☆
Arteriography abdomen	May Be Appropriate (Disagreement)	☆☆☆
MRA abdomen and pelvis without and with IV contrast	May Be Appropriate (Disagreement)	○
X-ray abdomen	May Be Appropriate	☆☆
US duplex Doppler abdomen	May Be Appropriate	○
CT abdomen and pelvis without and with IV contrast	Usually Not Appropriate	☆☆☆☆
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	☆☆☆☆
MRA abdomen and pelvis without IV contrast	Usually Not Appropriate	○
<b>Variant 2:</b> Suspected chronic mesenteric ischemia. Initial imaging.		
Procedure	Appropriateness Category	Relative Radiation Level
CTA abdomen and pelvis with IV contrast	Usually Appropriate	☆☆☆
MRA abdomen and pelvis without and with IV contrast	Usually Appropriate	○
Arteriography abdomen	May Be Appropriate (Disagreement)	☆☆☆
CT abdomen and pelvis with IV contrast	May Be Appropriate	☆☆☆
MRA abdomen and pelvis without IV contrast	May Be Appropriate	○
US duplex Doppler abdomen	May Be Appropriate	○
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	☆☆☆
CT abdomen and pelvis without and with IV contrast	Usually Not Appropriate	☆☆☆☆
X-ray abdomen	Usually Not Appropriate	☆☆

87 yo woman with HTN, CAD, diabetes, dementia presents with severe and acute abdominal pain. Abdominal pain is maybe ?out of proportion to physical examination.

Now what? Do I order a study?  
What study do I order?



## Part V: Oral and Intravenous Contrast for CT Examinations

### Evaluation of Acute Abdominal Pain in the Emergency Setting Using Computed Tomography Without Oral Contrast in Patients With Body Mass Index Greater Than 25.

[Uyeda JW<sup>1</sup>](#), [Yu H](#), [Ramalingam V](#), [Devalapalli AP](#), [Soto JA](#), [Anderson SW](#).

#### ⊕ Author information



#### Abstract

**PURPOSE:** To evaluate the rate of delayed or missed diagnoses and need for additional computed tomography (CT) imaging in emergency department patients with abdominal pain who are imaged without oral contrast.

**MATERIALS AND METHODS:** The institutional review board approved this Health Insurance Portability and Accountability Act-compliant retrospective study; informed consent was waived. All consecutive adult patients with body mass index greater than 25 undergoing a CT abdomen/pelvis with intravenous contrast and without oral contrast with nontraumatic acute abdominal pain during a 16-month period at our academic tertiary care center were included. Medical records were reviewed, imaging findings on admission CT, use of repeat CT examinations within 4 weeks of the original examination, and clinical outcomes were recorded. In patients undergoing repeat imaging, an investigator determined whether repeat imaging was influenced by the lack of oral contrast on the original examination. As the most common cause of bowel-related positive CT scans, an analysis of acute appendicitis was performed.

**RESULTS:** Of the 1992 patients included in this study, 4 patients (0.2%) underwent repeat CT studies directly related to the absence of oral contrast on the original examination. Of the 1992 CT scans, 1193(59.8%) were interpreted as negative, none of which required surgery or direct intervention. In patients with acute appendicitis, there was a sensitivity of CT in this patient population of 100% with a specificity of 99.5%.

**CONCLUSIONS:** In patients with body mass index greater than 25 presenting to the ED with acute abdominal pain, CT examinations can be acquired without oral contrast without compromising the clinical efficacy of CT.







## Use of PO Contrast

- Recent postoperative patients (~30 days)
- Penetrating trauma (stab wound, GSW)
- At the request of the surgeon


## Utility of IV Contrast

3/4/2017 6:00 AM



No stones or hydronephrosis


3/4/2017 11:35 PM




Renal infarct


S/P Appendectomy

+IV  
Contrast





Anastomotic  
& abdominal  
wall  
abscesses





Soft Tissue Hematoma



+IV  
Contrast




Pseudoaneurysm in  
soft tissue hematoma

Requiring coil  
embolization



## Abdominal Pain







## Nephrolithiasis Noncontrast CT






## Nephroureterolithiasis +IV CT (Routine PV phase)




R proximal ureteral stone w hydronephrosis, delayed nephrogram



## Uric Acid Stones analysis by Dual Energy CT



## Part VI: Renal Failure and Contrast Allergies



www.acr.org/Clinical-Resources/Contrast-Manual

### Contrast-Induced Nephropathy

- “At the current time, it is the position of ACR Committee on Drugs and Contrast Media that CIN is a real, albeit rare, entity.”
- Literature fails to include control group
- Studies done on cardiac angiography patients → overestimate CIN risk
- Threshold **eGFR  $\geq 30$  mL/min/1.73m<sup>2</sup>**

## Risk Factors Warranting Renal Function Assessment

- Age > 60
- History of renal disease, including:
  - Dialysis
  - Renal cancer
  - Kidney transplant
  - Single kidney
  - Renal surgery
- History of hypertension requiring medical therapy
- History of diabetes mellitus
- Metformin or metformin-containing drug combinations

\*Patients scheduled for routine intravascular study but do NOT have one of the above risk factors do NOT require a baseline serum creatinine determination before iodinated contrast medium administration.

[www.acr.org/Clinical-Resources/Contrast-Manual](http://www.acr.org/Clinical-Resources/Contrast-Manual)

## Dialysis Patients

- Patients with anuric end-stage renal disease who do not have functioning transplant can receive IV contrast
- Theoretical risk of converting oliguric patient on dialysis to anuric patient but remains speculative
- Low-osmolality contrast is readily cleared by dialysis. Unless unusually large volume of contrast is given, or there is substantial underlying cardiac dysfunction, there is **NO** need for urgent dialysis after contrast

[www.acr.org/Clinical-Resources/Contrast-Manual](http://www.acr.org/Clinical-Resources/Contrast-Manual)



## Allergic Reactions

- Prednisone-based:
  - 50 mg prednisone by mouth 13, 7, and 1 hour before contrast administration + 50 mg diphenhydramine IV, IM, or PO 1 hour before contrast administration
- OR
- Methylprednisolone-based:
  - 32 mg methylprednisolone by mouth 12 hours and 2 hours before contrast administration. 50 mg diphenhydramine may be added as an option.

[www.acr.org/Clinical-Resources/Contrast-Manual](http://www.acr.org/Clinical-Resources/Contrast-Manual)


## Allergic Reactions

- Premedication does not prevent all reactions
- Has not been confirmed to reduce moderate to severe reactions or reaction-related deaths
- Limited supporting efficacy in high-risk patients
- History of prior severe contrast reaction is considered a **relative contraindication**
- Rare situations where urgency of contrast study may outweigh benefits of prophylaxis → must be made jointly by radiology, referring service, and the patient (if feasible) and resuscitation team should be available

[www.acr.org/Clinical-Resources/Contrast-Manual](http://www.acr.org/Clinical-Resources/Contrast-Manual)

## Lecture Summary

- Interpret basic chest radiographs
- Recognize and locate various tubes and lines
- Identify “can’t miss radiology diagnoses” on plain x-ray and CT
- Discuss various imaging protocols and considerations
- Recognize the pros and cons of IV and PO contrast use in CT
- Optimally triage renal failure and contrast allergy patients



Thank you!

[juyeda@bwh.harvard.edu](mailto:juyeda@bwh.harvard.edu)

