

























































































No Bridging Outcome Bridging (N-93)8 Prval Prumber of patients (percent) Prval Primary Primary
number of patients (percent) Primary Arterial thromboembolism 4 (0.4) 3 (0.3) 0.01°, 0 Stroke 2 (0.2) 3 (0.3)
Primary Arterial thromboembolism 4 (0.4) 3 (0.3) 0.01°, 0 Stroke 2 (0.2) 3 (0.3)
Arterial thromboembolism 4 (0.4) 3 (0.3) 0.01°, 0 Stroke 2 (0.2) 3 (0.3)
Stroke 2 (0.2) 3 (0.3)
Transient ischemic attack 2 (0.2) 0
Systemic embolism 0 0
Major bleeding 12 (1.3) 29 (3.2) 0.00
Secondary
Death 5 (0.5) 4 (0.4) 0.88
Myocardial infarction 7 (0.8) 14 (1.6) 0.10
Deep-vein thrombosis 0 1 (0.1) 0.25
Pulmonary embolism 0 1 (0.1) 0.25
Minor bleeding 110 (12.0) 187 (20.9) <0.00

Bridging Anticoagulation		
CHADS ₂ score‡	No Bridging (N = 950)	Bridging (N = 934)
Mean	2.3±1.03	2.4±1.07
Distribution — no. (%)		
0	1 (0.1)	1 (0.1)
1	216 (22.7)	212 (22.7)
2	382 (40.2)	351 (37.6)
3	229 (24.1)	232 (24.8)
4	96 (10.1)	106 (11.3)
5	23 (2.4)	27 (2.9)
6	3 (0.3)	5 (0.5)
IGUIGE BOUREIS JO, Spyropoulos AC, Radio S, et al Perforenze de Régiong Actionogulation in Potentis with Antial Phillipsion ACMA. 2015;373(9):823-833.		



























