



HARVARD MEDICAL SCHOOL
TEACHING HOSPITAL



Mass General Brigham

Best Practices for Inpatient Management of Patients with Psychiatric Illness

Sejal B. Shah, MD, FACLP, DFAPA
Chief, Division of Medical Psychiatry
Associate Vice Chair, Clinical Consultation Services
Brigham and Women's Hospital
Assistant Professor, Harvard Medical School

Disclosure

With respect to the following presentation, there has been no relevant (direct or indirect) financial relationship with any for-profit company which could be considered a conflict of interest.

Outline

- Demoralization
- Depression
- Anxiety
- Somatic Symptom Disorder
- Insomnia
- Personality Disorders
- Trauma-Informed Care
- Psychotic Disorders
- Take Home Points

Fiorillo and Sartorius
Annals of General Psychiatry (2021) 20:52
<https://doi.org/10.1186/s12991-021-00374-y>

Annals of General Psychiatry

COMMENTARY

Open Access

Mortality gap and physical comorbidity of people with severe mental disorders: the public health scandal



Andrea Fiorillo^{1*} and Norman Sartorius²

Conclusion: “Patients suffering from severe mental disorders, including schizophrenia, major depression and bipolar disorders, have a reduced life expectancy compared to the general population of up to 10–25 years.”

Case Example

47 yo man with h/o DM2, HTN, hyperlipidemia, asthma and no formal past psychiatric history is admitted to the hospital with acute renal failure. His hospitalization is complicated by PNA followed by sepsis. He is admitted to the ICU, intubated and has a prolonged stay of 1 month. He is extubated and learns that he will need lifelong dialysis as he is not a candidate for renal transplantation. His medical team notes that he sleeps most of the day, does not engage in physical therapy and eats very little. He notes that he has periods of the day when he is in intense pain and does not know how he will learn to live with this new obstacle in his life. He denies depressed mood and his cognitive status remains intact. His mood is visibly bright when his children visit.

Conservation-Withdrawal Versus Depression

Myron F. Weiner, M.D.

Clinical Professor of Psychiatry

Robert Lovitt, Ph.D.

Associate Professor of Psychiatry

University of Texas Health Science Center at Dallas, Dallas, Texas

General Hospital Psychiatry
© Elsevier North Holland, Inc., 1979



Demoralization

- No Diagnostic Criteria
- Definition: dysphoric state, destruction of confidence/hope, disempowerment
- Contrasted with Depression
- Prevalence: 1/3 of physically ill patients

Management of Demoralization

Encouragement,
Enhanced
Engagement

Validate,
Normalize
Experience of
Illness

Active Symptom
Management

Exploration of
Attitudes of Hope
and Meaning in
Life

Searching for
Renewed Purpose
and Role in Life

Cognitive
Restructuring of
Negative Beliefs

Spiritual/Religious
Support

Promotion of
Social Contact

Case Example

26 yo woman with h/o depression was admitted s/p MVA with resultant concussion, multiple rib fractures and tibia fracture. She is now post-op ORIF of tibia. She reports to you that she has not been taking her antidepressant for many months now and that this accident has led to an overall feeling of hopelessness. She cites that her appetite is low, it's difficult for her to concentrate and "I just don't know if I have the energy to do all this."

Depression Symptoms

5 or More Symptoms During 2-Week Period

- At least 1 of:
 - Depressed Mood
 - Loss of Interest or Pleasure (Anhedonia)
- Other Symptoms:
 - Significant Weight Loss or Decrease or Increase in Appetite Nearly Every Day
 - Thought Slowing/Reduction of Physical Movement
 - Fatigue or Loss of Energy Nearly Every Day
 - Feelings of Worthlessness or Inappropriate Guilt
 - Decreased Ability to Concentrate
 - Recurrent Thoughts of Death/Recurrent Suicidal Ideation



Depression

- Consequences of Depression
 - Poorer Physical Health
 - Worse Functional Outcomes
 - Greater Rates of Rehospitalization
- Prevalence
- Overlap with Demoralization
- Overlap with Delirium

Suicidality

The Joint Commission Journal on Quality and Patient Safety 2018; 44:643–650

Incidence and Method of Suicide in Hospitals in the United States

Scott C. Williams, PsyD; Stephen P. Schmalz, PhD; Genard M. Castro, PhD, MPH; David W. Baker, MD, MPH

Psychiatric Safety Risk Assessment

- Active vs Passive Suicidal Ideation
- Risk Factors
- Protective Factors
- Validated Screening Tools
- Collateral Information

Case Example

45 yo man with no previous medical history was hospitalized x 1.5 months s/p fall from ladder leading to SCI. The patient did well during his hospitalization, participating in the treatment plan with motivation. Close to the time of discharge, the patient began to show reticence towards participation in discharge planning and his mood seemed to have shifted towards irritability and extreme worry.

Anxiety Disorders

- Separation Anxiety Disorder
- Specific Phobias
- Social Anxiety Disorder
- Panic Disorder
- Generalized Anxiety Disorder
- Substance/Medication-Induced Anxiety Disorder

Anxiety

- Excessive worry occurring more days than not for at least 6 months about a number of events or activities
- Difficult to control the worry
- 3 or More:
 - Restlessness/Feeling on Edge
 - Easily Fatigued
 - Difficulty Concentrating
 - Irritability
 - Muscle Tension
 - Sleep Disturbance

Anxiety: Barrier to Care

- Previous History of Anxiety?
- Explore Specific Fears
- Transparency of Treatment Plan
 - Loss of Control
- Avoid Reassurance or “Advice Mode”
- Referral to Cognitive Behavioral Therapy (CBT)
- Medications

Case Example

45 yo woman with h/o anxiety, DM2 was admitted to the hospital with chest and abdominal pain and slightly elevated 1st troponin. Remainder of the cardiac and abdominal workup is unrevealing, but patient remains with pain. She reports that's she is convinced “there's something really wrong” and cannot be reassured. She begins demanding unnecessary studies and discharge planning is difficult.

Somatic Symptom Disorder

- One or more somatic symptoms that are distressing or result in significant disruption of daily life
- Excessive thoughts, feelings or behaviors related to the somatic symptoms or associated health concerns as manifested by:
 - Disproportionate and persistent thoughts about the seriousness of one's symptoms
 - Persistently high level of anxiety about health or symptoms
 - Excessive time and energy devoted to these symptoms or health concerns
- Although any one somatic symptom may not be continuously present, the state of being symptomatic is persistent (typically more than 6 months)

Somatic Symptom Disorder: Management

Encourage Stable and Consistent Primary Care

Support/Understanding

Validation of Fears

Suggest Tracking Symptom Information

Encourage Family/Friend Involvement

Management of Underlying Psychiatric Disorders

Insomnia

- Behavioral Interventions/Sleep Hygiene
 - Fixed Sleep/Wake Time
 - Minimize Daytime Sleeping
 - Unplug from Electronics
 - Relaxation Techniques
 - Quiet Environment/Minimize Disturbances
- Underlying Conditions
 - Anxiety
 - Depression
 - Delirium
 - Pain
 - Psychosocial Stressors
- Medication Management

Case Example

39 yo woman with h/o borderline personality disorder, depression, HTN, CHF who was admitted for CHF exacerbation. The medical team notes that her chart indicates previous “demanding behaviors.” During this admission, the patient is disparaging of staff, demanding in her behaviors and intermittently refuses to participate in care. She makes comments that some providers are “the best” and refuses to work with others.

Personality Disorders

Borderline Personality Disorder

- Efforts to Avoid Real or Imagined Abandonment
- Pattern of Unstable and Intense Interpersonal Relationships
 - Alternates Between Extremes of Idealization and Devaluation
- Identity Disturbance
- Impulsivity in 2 Areas (spending, sex, substance use, reckless driving, binge eating)
- Recurrent Suicidal Behavior, Gestures/Threats, Self-Mutilating Behavior
- Affective Instability
- Chronic Feelings of Emptiness
- Inappropriate, Intense Anger
- Transient, Stress-Related Paranoid Ideation or Severe Dissociative Symptoms

Personality Disorders-- Management

Avoid Personalization

- Monitor Countertransference
- Tolerate Intense Anger or Hate

Set Limits

Consistent Messaging to Avoid Splitting

Reinforce Positive Behavior

Medications?

Borderline Personality Disorder A Review

Falk Leichsenring, DSc; Nikolas Heim, MA, MSc; Frank Leweke, MD; Carsten Spitzer, MD;
Christiane Steinert, PhD; Otto F. Kernberg, MD

Leichsenring F, Heim N, Leweke F, Spitzer C, Steinert C, Kernberg OF. Borderline Personality Disorder: A Review. JAMA. 2023 Feb 28;329(8):670-679. PMID: 36853245.

Trauma

- Types of Trauma
- Prevalence
- Poorer Health Outcomes
- Shame and Stigma
- Risk of Re-Traumatization
- Adverse Childhood Experiences (ACEs)

Trauma-Informed Care: Guiding Principles



Case Example

54 yo man with h/o schizoaffective disorder-bipolar type, DM, HTN, Crohn's disease presented to the hospital for scheduled bowel resection surgery. Post-surgery, the patient is initially calm but then begins to exhibit paranoia towards staff (worried that nurses are talking about him, refuses to eat hospital food) and intermittent refusal to participate in care. The patient's cognition is intact and delirium is ruled out.

Psychotic Disorders

- Psychotic Disorders
 - Delusional Disorder
 - Schizophrenia
 - Schizoaffective Disorder
 - Substance/Medication-Induced Psychotic Disorder
 - Psychotic Disorder Due to a Medical Condition
- (Bipolar Disorder)
- Hallmark Symptoms
 - Delusions
 - Hallucinations
 - Disorganized Thinking and/or Behavior
 - Negative Symptoms

Case Example— Revisited

54 yo man with h/o schizoaffective disorder-bipolar type, DM, HTN, Crohn's disease presented to the hospital for scheduled bowel resection surgery. Post-surgery, the patient is initially calm but then begins to exhibit paranoia towards staff (worried that nurses are talking about him, refuses to eat hospital food) and intermittent refusal to participate in care. The patient's cognition is intact and delirium is ruled out.

- Reassure Safety
- Do Not Challenge Delusions
- Transparency of Interventions and Treatment Plan
- Build Rapport
- Medication Management

Take Home Points

- Demoralization vs Depression
- Recognition of Suicidality
- Consequences of Anxiety
- Reassurance and Validation of Patients with Somatic Symptom Disorder
- Importance of Insomnia Management in the Inpatient Setting
- Management of Personality Disorders
- Psychosis Can Be Managed Effectively in a Hospital Setting